

Medidas de prevención y control (o erradicación) de la salmonelosis

Maurice Raccoursier

H&N Internacional

Referencias bibliográficas

Bailey, J. S., Cox, N. A., & Berrang, M. E. (1994). Hatchery-acquired salmonellae in broiler chicks. *Poult Sci*, 73(7), 1153-1157. <https://doi.org/10.3382/ps.0731153>

Bailey, J. S., Cox, N. A., Blankenship, L. C. & Stern, N. J. (1992). Hatchery contamination reduces the effectiveness of competitive exclusion treatments to control salmonella colonization of broiler chicks. *Poultry Sci*. 71(Suppl. 1):6 (Abstr).

Cameron, D. M., Carter, J. N., & Mansell, P. (1996). Evaluation of Aviguard against a Salmonella enteritidis infective model in broiler chickens. In *Western Poultry Disease Conference (USA)*.

Davies, R. H., & Wray, C. (1995). Mice as Carriers of Salmonella-Enteritidis on Persistently Infected Poultry Units. *Veterinary Record*, 137(14), 337-341. <https://doi.org/DOI 10.1136/vr.137.14.337>

Davies, R. H., & Wray, C. (1995). Mice as Carriers of Salmonella-Enteritidis on Persistently Infected Poultry Units. *Veterinary Record*, 137(14), 337-341. <https://doi.org/DOI 10.1136/vr.137.14.337>

Ekperigin, H.E., McCapes, R.H. & Redus, R., Ritchie, W.L., & Cameron, W.J. (1991). Production of poultry feed free of salmonella and *Escherichia coli*. 10.1016/B978-0-12-104280-6.50033-2.

Fernandez, F., Hinton, M., & Van Gils, B. (2002). Dietary mannan-oligosaccharides and their effect on chicken caecal microflora in relation to Salmonella Enteritidis colonization. *Avian Pathology*, 31(1), 49-58. <https://doi.org/Doi 10.1080/03079450120106000>

Gast, R. K., & Beard, C. W. (1992). Detection and Enumeration of Salmonella enteritidis in Fresh and Stored Eggs Laid by Experimentally Infected Hens. *J Food Prot*, 55(3), 152-156. <https://doi.org/10.4315/0362-028X-55.3.152>

Hofacre, C.L. (2023). *Salmonella* and salmonellosis. POPH 7060 lecture. MAM program. Poultry Research and Diagnostic Center, University of Georgia.

- Himathongkham, S., Pereira, M. G., & Riemann, H. (1996). Heat destruction of Salmonella in poultry feed: effect of time, temperature, and moisture. *Avian Dis*, 40(1), 72-77.
<https://www.ncbi.nlm.nih.gov/pubmed/8713051>
- Jones, F. T., Axtell, R. C., Rives, D. V., Scheideler, S. E., Tarver, F. R., Walker, R. L., & Wineland, M. J. (1991). A Survey of Salmonella Contamination in Modern Broiler Production. *Journal of Food Protection*, 54(7), 502-507. <https://doi.org/Doi.10.4315/0362-028x-54.7.502>
- Jones, P. W., Dougan, G., Hayward, C., Mackenzie, N., Collins, P., & Chatfield, S. N. (1991). Oral Vaccination of Calves against Experimental Salmonellosis Using a Double Aro Mutant of Salmonella-Typhimurium. *Vaccine*, 9(1), 29-34. [https://doi.org/Doi.10.1016/0264-410x\(91\)90313-U](https://doi.org/Doi.10.1016/0264-410x(91)90313-U)
- Koutsoumanis, K., Allende, A., Alvarez-Ordóñez, A., Bolton, D., Bover-Cid, S., Chemaly, M, De Cesare, A., Herman, L., Hilbert, F., Lindqvist, R., Nauta, M., Peixe, L., Ru, G., Simmons, M., Skandamis, P., Suffredini, E., Dewulf, J., Hald, T., Michel, V. N., Ricci, A., Snary, E., Boelaert, F., Messens, W., and Davies, R. (2019). Scientific Opinion on the *Salmonella* control in poultry flocks and its public health impact. *EFSA Journal* 2019;17(2):5596, 155 pp. <https://doi.org/10.2903/j.efsa.2019.5596>
- Kim, A., Lee, Y. J., Kang, M. S., Kwag, S. I., & Cho, J. K. (2007). Dissemination and tracking of Salmonella spp. in integrated broiler operation. *Journal of Veterinary Science*, 8(2), 155-161.
<https://doi.org/DOI.10.4142/jvs.2007.8.2.155>
- Liu, T. S., Snoeyenbos, G. H., & Carlson, V. L. (1969). The effect of moisture and storage temperature on a Salmonella senftenberg 775W population in meat and bone meal. *Poult Sci*, 48(5), 1628-1633.
<https://doi.org/10.3382/ps.0481628>
- Matlho, G., Himathongkham, S., Riemann, H., & Kass, P. (1997). Destruction of Salmonella enteritidis in poultry feed by combination of heat and propionic acid. *Avian Dis*, 41(1), 58-61.
<https://www.ncbi.nlm.nih.gov/pubmed/9087320>
- Matlho, G., Himathongkham, S., Riemann, H., & Kass, P. (1997). Destruction of Salmonella enteritidis in poultry feed by combination of heat and propionic acid. *Avian Diseases*, 41(1), 58-61.
<https://doi.org/Doi.10.2307/1592443>
- Northcutt, J. K., Savage, S. I., & Vest, L. R. (1997). Relationship between feed withdrawal and viscera condition of broilers. *Poultry Science*, 76(2), 410-414. <https://doi.org/DOI.10.1093/ps/76.2.410>

- Oyarzabal, O. A., & Conner, D. E. (1996). Application of direct-fed microbial bacteria and fructooligosaccharides for salmonella control in broilers during feed withdrawal. *Poult Sci*, 75(2), 186-190. <https://doi.org/10.3382/ps.0750186>
- Parker, E. M., Parker, A. J., Short, G., O'Connor, A. M., & Wittum, T. E. (2022). Salmonella detection in commercially prepared livestock feed and the raw ingredients and equipment used to manufacture the feed: A systematic review and meta-analysis. *Preventive Veterinary Medicine*, 198. <https://doi.org/ARTN 10554610.1016/j.prevetmed.2021.105546>
- Ramirez, G. A., Sarlin, L. L., Caldwell, D. J., Yezak, C. R., Jr., Hume, M. E., Corrier, D. E., Deloach, J. R., & Hargis, B. M. (1997). Effect of feed withdrawal on the incidence of Salmonella in the crops and ceca of market age broiler chickens. *Poult Sci*, 76(4), 654-656. <https://doi.org/10.1093/ps/76.4.654>
- Rigby, C. E., & Pettit, J. R. (1981). Effects of feed withdrawal on the weight, fecal excretion and Salmonella status of market age broiler chickens. *Can J Comp Med*, 45(4), 363-365. <https://www.ncbi.nlm.nih.gov/pubmed/7039784>
- Roche, A. J., Cox, N. A., Richardson, L. J., Buhr, R. J., Cason, J. A., Fairchild, B. D., & Hinkle, N. C. (2009). Transmission of Salmonella to broilers by contaminated larval and adult lesser mealworms, *Alphitobius diaperinus* (Coleoptera: Tenebrionidae). *Poultry Science*, 88(1), 44-48. <https://doi.org/10.3382/ps.2008-00235>
- Rolon. (2005). Salmonella vaccination studies in breeders and their progeny: relating humoral and mucosal immunity with resistance to challenge [University of Georgia]. http://getd.libs.uga.edu/pdfs/rolon_ariel_200505_phd.pdf
- Schneitz, C., Koivunen, E., Tuunainen, P., & Valaja, J. (2016). The effects of a competitive exclusion product and two probiotics on Salmonella colonization and nutrient digestibility in broiler chickens. *Journal of Applied Poultry Research*, 25(3), 396-406. <https://doi.org/10.3382/japr/pfw025>
- Skov, M. N., Spencer, A. G., Hald, B., Petersen, L., Nauerby, B., Carstensen, B., & Madsen, M. (2004). The role of litter beetles as potential reservoir for Salmonella enterica and thermophilic Campylobacter spp. between broiler flocks. *Avian Diseases*, 48(1), 9-18. <https://doi.org/Doi 10.1637/5698>
- Suganuma, K., Hamasaki, T., & Hamaoka, T. (2021). Effect of dietary direct-fed microbial and yeast cell walls on cecal digesta microbiota of layer chicks inoculated with nalidixic acid resistant Salmonella Enteritidis. *Poult Sci*, 100(10), 101385. <https://doi.org/10.1016/j.psj.2021.101385>

- Wang, J., Vaddu, S., Bhumanapalli, S., Mishra, A., Applegate, T., Singh, M., & Thippareddi, H. (2023). A systematic review and meta-analysis of the sources of Salmonella in poultry production (pre-harvest) and their relative contributions to the microbial risk of poultry meat. *Poultry Science*, 102(5). [https://doi.org/ARTN 102566](https://doi.org/ARTN%20102566) 10.1016/j.psj.2023.102566
- Weinack, O. M., Snoeyenbos, G. H., & Soerjadiliem, A. S. (1985). Further-Studies on Competitive-Exclusion of Salmonella-Typhimurium by Lactobacilli in Chickens. *Avian Diseases*, 29(4), 1273-1276. [https://doi.org/Doi 10.2307/1590485](https://doi.org/Doi%2010.2307/1590485)